

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁵ : A61K 37/28</p>	<p>A1</p>	<p>(11) International Publication Number: WO 93/18786 (43) International Publication Date: 30 September 1993 (30.09.93)</p>
<p>(21) International Application Number: PCT/DK93/00099 (22) International Filing Date: 18 March 1993 (18.03.93) (30) Priority data: 363/92 19 March 1992 (19.03.92) DK (71) Applicant (for all designated States except US): NOVO NORDISK A/S [DK/DK]; Novo Allé, DK-2880 Bagsvaerd (DK). (71)(72) Applicants and Inventors: EFENDIC, Suad [SE/SE]; Stjärnvägen 16B, S-181 34 Lidingö (SE). GUTNIAK, Mark [SE/SE]; Hässelby Strandväg 26, S-165 65 Hässelby (SE).</p>		<p>(72) Inventor; and (75) Inventor/Applicant (for US only) : KIRK, Ole [DK/DK]; Stefansgade 38, 3/tv., DK-2200 Copenhagen N (DK). (74) Common Representative: NOVO NORDISK A/S; Patent Department, Novo Allé, DK-2880 Bagsvaerd (DK). (81) Designated States: AU, BB, BG, BR, CA, CZ, FI, HU, JP, KP, KR, KZ, LK, MG, MN, MW, NO, NZ, PL, RO, RU, SD, SK, UA, US, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, SN, TD, TG). Published With international search report.</p>
<p>(54) Title: USE OF A PEPTIDE</p> <p>(57) Abstract</p> <p>GLP-1(7-37), GLP-1(7-36)amide, and certain related compounds can be used to provide a medicament for use in the treatment of diabetes in a regimen which additionally comprises treatment with an oral hypoglycaemic agent. A strong synergistic effect is observed.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NL	Netherlands
BE	Belgium	GN	Guinea	NO	Norway
BF	Burkina Faso	GR	Greece	NZ	New Zealand
BG	Bulgaria	HU	Hungary	PL	Poland
BJ	Benin	IE	Ireland	PT	Portugal
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SK	Slovak Republic
CI	Côte d'Ivoire	LJ	Liechtenstein	SN	Senegal
CM	Cameroun	LK	Sri Lanka	SU	Soviet Union
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	MC	Monaco	TC	Togo
DE	Germany	MG	Madagascar	UA	Ukraine
DK	Denmark	ML	Mali	US	United States of America
ES	Spain	MN	Mongolia	VN	Viet Nam
FI	Finland				

USE OF A PEPTIDE

FIELD OF THE INVENTION

The present invention relates to the use of GLP-1(7-37), GLP-1(7-36)amide, or certain related compounds
5 for the preparation of a medicament for use in the treatment of diabetes in a regimen which additionally comprises treatment with an oral hypoglycaemic agent. The invention also relates to a method of treating diabetes by using said medicament.

10 BACKGROUND OF THE INVENTION

Diabetes is characterized by an impaired glucose metabolism manifesting itself among other things by an elevated blood glucose level in the diabetic patients. Underlying defects lead to a classification of diabetes into
15 two major groups: type 1 diabetes, or insulin demanding diabetes mellitus (IDDM), which arises when patients lack β -cells producing insulin in their pancreatic glands, and type 2 diabetes, or non-insulin dependent diabetes mellitus (NIDDM), which occurs in patients with an impaired β -cell
20 function besides a range of other abnormalities.

Type 1 diabetic patients are currently treated with insulin, while the majority of type 2 diabetic patients are treated either with agents that stimulate β -cell function or with agents that enhance the tissue sensitivity of
25 the patients towards insulin.

Among the agents applied for stimulation of the β -cell function, those acting on the ATP-dependent potassium channel of β -cells are most widely used in current therapy. The so-called sulfonylureas such as tolbutamide,
30 glibenclamide, glipizide, and gliclazide are used extensively and other agents such as AG-EE 623 ZW also acting at this

molecular site are under development (AG-EE 623 ZW is a company code for (S)-(+)-2-ethoxy-4-[2-[[3-methyl-1-[2-(1-piperidinyl)phenyl]butyl]-amino]-2-oxoethyl]benzoic acid, a compound described in European patent publication No. 5 147,850 (to Dr. Karl Thomae GmbH)). Among the agents applied to enhance tissue sensitivity towards insulin metformin is a representative example.

Even though sulfonylureas are widely used in the treatment of NIDDM this therapy is, in most instances, not 10 satisfactory: In a large number of NIDDM patients sulfonylureas do not suffice to normalize blood sugar levels and the patients are, therefore, at high risk for acquiring diabetic complications. Also, many patients gradually lose the ability to respond to treatment with sulfonylureas and are thus 15 gradually forced into insulin treatment. This shift of patients from oral hypoglycaemic agents to insulin therapy is usually ascribed to exhaustion of the β -cells in NIDDM patients.

Over the years, numerous attempts have therefore 20 been made to provide novel agents which stimulate β -cell function in order to offer the NIDDM patients an improved treatment. Recently, a series of peptides derived from glucagon-like peptide-1 have been considered as insulino-tropic agents for therapeutic use.

25 Glucagon-like peptide-1, also referred to as GLP-1, is a peptide sequence found in the C-terminal portion of mammalian proglucagon. Prior to 1985, no definite biological activity of GLP-1 had been reported. However, in 1985 it was demonstrated that the amide of a fragment of GLP-1, 30 namely GLP-1(1-36)amide, stimulates insulin release from isolated precultured rat pancreatic islets in the presence of glucose in a dose-dependent manner (Schmidt, W.E. et al. Diabetologia 28 (1985) 704-7). This finding suggests that GLP-1(1-36)amide and related peptides might be useful in the 35 treatment of type 2 diabetes. Due to its substantially closer sequence homology to glucagon and glucosedependent insulinotropic peptide, also referred to as GIP, Schmidt et

al. suggested that an even stronger glucagon- and/or GIP-like biological activity could be expected with GLP-1(7-36) than with the intact peptide. In recent years, particular interest has focused on the GLP-1 fragments GLP-1(7-37) and
5 GLP-1(7-36)amide and analogues and functional derivatives thereof. The designation GLP-1(1-36) indicates that the peptide fragment in question comprises the amino acid residues from (and including) number 1 to (and including) number 36 when counted from the N-terminal end of the parent
10 peptide, GLP-1. Similarly, the designation GLP-1(7-37) designates that the fragment in question comprises the amino acid residues from (and including) number 7 to (and including) number 37 when counted from the N-terminal end of the parent peptide, GLP-1. The amino acid sequence of
15 GLP-1(7-36)amide and of GLP-1(7-37) is given in formula I:

His-Ala-Glu-Gly-Thr-Phe-Thr-Ser-Asp-Val-Ser-
Ser-Tyr-Leu-Glu-Gly-Gln-Ala-Ala-Lys-Glu-Phe-
Ile-Ala-Trp-Leu-Val-Lys-Gly-Arg-X

(I)

20 which shows GLP-1(7-36)amide when X is NH₂ and GLP-1(7-37) when X is Gly-OH.

That GLP-1(7-36)amide is indeed an insulino-tropic agent in man has been demonstrated by Kreymann, B. et al. who infused this peptide into healthy volunteers and
25 observed a significant rise in plasma insulin (Lancet 2 (1987) 1300-304).

The insulintropic action of GLP-1(7-37) in diabetic as well as in nondiabetic subjects has been demonstrated by Nathan, D.M. et al. Diabetes Care 15 (1992) 270-
30 76.

International Patent Application No. WO 87/06941 (to The General Hospital Corporation) relates to a peptide fragment which comprises GLP-1(7-37) and functional derivatives thereof and to its use as an insulintropic agent.

35 International Patent Application No. 90/11296

(to The General Hospital Corporation) relates to a peptide fragment which comprises GLP-1(7-36) and functional derivatives thereof and has an insulintropic activity which exceeds the insulintropic activity of GLP-1(1-36) or GLP-1
5 (1-37) and to its use as an insulintropic agent.

International Patent Application No. 91/11457 (to Buckley et al.) relates to effective analogs of the active GLP-1 peptides 7-34, 7-35, 7-36, and 7-37.

The effect of GLP-1(7-37) in combination with
10 glibenclamide on insulin secretion from rat pancreatic islets was studied in vitro by Parker, J.C. et al. (Diabetes 40 (suppl. 1) (1991) 237 A). Only an additive effect of the two agents was observed.

However, to the best of the knowledge of the
15 present inventors the surprising synergistic effect in vivo achieved by the combined use of an oral hypoglycaemic agent and a fragment of GLP-1 or an analogue or a functional derivative thereof has not previously been disclosed.

SUMMARY OF THE INVENTION

20 The present invention relates to the surprising finding that when GLP-1 related peptides are administered in combination with oral hypoglycaemic agents in general and with sulfonylureas in particular for treatment of type 2 diabetes, a synergistic effect is observed. This surprising
25 observation has been made even in type 2 diabetic patients who fail to respond when sulfonylureas are administered alone.

Thus, in its broadest aspect the present invention relates to the use of GLP-1(7-27), GLP-1(7-36)amide, or
30 a pharmaceutically acceptable peptide containing a fragment of the GLP-1(7-37) sequence, or an analogue or a functional derivative of such a peptide for the preparation of a medicament for use in the treatment of type 2 diabetes in a regimen which additionally comprises treatment with an oral

hypoglycaemic agent and to a method of treating type 2 diabetes which method comprises administering an effective amount of GLP-1(7-37), GLP-1(7-36)amide, or a pharmaceutically acceptable peptide containing a fragment of the
5 GLP-1(7-37) sequence, or an analogue or a functional derivative of such a peptide to a patient in a regimen which additionally comprises treatment with an oral hypoglycaemic agent.

In a first preferred embodiment, the present in-
10 vention relates to the use of GLP-1(7-36)amide for the preparation of a medicament for use in the treatment of type 2 diabetes in a regimen which additionally comprises treatment with an oral hypoglycaemic agent.

In a further preferred embodiment, the present
15 invention relates to the use of GLP-1(7-37) for the preparation of a medicament for use in the treatment of type 2 diabetes in a regimen which additionally comprises treatment with an oral hypoglycaemic agent.

In a further preferred embodiment, the present
20 invention relates to the use of an analogue of GLP-1(7-37) for the preparation of a medicament for use in the treatment of type 2 diabetes in a regimen which additionally comprises treatment with an oral hypoglycaemic agent.

In a further preferred embodiment, the present
25 invention relates to the use of a functional derivative of GLP-1(7-37) for the preparation of a medicament for use in the treatment of type 2 diabetes in a regimen which additionally comprises treatment with an oral hypoglycaemic agent.

30 In a further preferred embodiment, the present invention relates to the use of GLP-1(7-37) or a fragment thereof or an analogue or a functional derivative of any of these including GLP-1(7-36)amide for the preparation of a medicament for use in the treatment of type 2 diabetes in a
35 regimen which additionally comprises treatment with tolbutamide.

In a further preferred embodiment, the present

invention relates to the use of GLP-1(7-37) or a fragment thereof or an analogue or a functional derivative of any of these including GLP-1(7-36)amide for the preparation of a medicament for use in the treatment of type 2 diabetes in a
5 regimen which additionally comprises treatment with glibenclamide.

In a further preferred embodiment, the present invention relates to the use of GLP-1(7-37) or a fragment thereof or an analogue or a functional derivative of any of
10 these including GLP-1(7-36)amide for the preparation of a medicament for use in the treatment of type 2 diabetes in a regimen which additionally comprises treatment with glipizide.

In a further preferred embodiment, the present
15 invention relates to the use of GLP-1(7-37) or a fragment thereof or an analogue or a functional derivative of any of these including GLP-1(7-36)amide for the preparation of a medicament for use in the treatment of type 2 diabetes in a regimen which additionally comprises treatment with
20 gliclazide.

In a further preferred embodiment, the present invention relates to the use of GLP-1(7-37) or a fragment thereof or an analogue or a functional derivative of any of these including GLP-1(7-36)amide for the preparation of a
25 medicament for use in the treatment of type 2 diabetes in a regimen which additionally comprises treatment with a biguanide.

In a further preferred embodiment, the present invention relates to the use of GLP-1(7-37) or a fragment
30 thereof or an analogue or a functional derivative of any of these including GLP-1(7-36)amide for the preparation of a medicament for use in the treatment of type 2 diabetes in a regimen which additionally comprises treatment with metformin.

35 In a further preferred embodiment, the present invention relates to the use of GLP-1(7-37) or a fragment

thereof or an analogue or a functional derivative of any of these including GLP-1(7-36)amide for the preparation of a medicament for use in the treatment of type 2 diabetes in a regimen which additionally comprises treatment with (S)-(+)-
5 2-ethoxy-4-[2-[[3-methyl-1-[2-(1-piperidinyl)phenyl]butyl]-amino]-2-oxoethyl]benzoic acid.

In this specification, analogues of GLP-1(7-37) or of GLP-1(7-36)amide, respectively, means peptides which differ from GLP-1(7-37) or from GLP-1(7-36)amide,
10 respectively, in that at least one of the amino acid residues of GLP-1(7-37) or of GLP-1(7-36)amide, respectively, independently has been exchanged by another amino acid residue, preferably one which can be coded for by the genetic code. The definition also comprises the case
15 when amino acid residues are added at or deleted from the N-terminal and/or the C-terminal end of the peptide. Preferably, the total number of such additions, deletions and exchanges does not exceed five, more preferred it does not exceed three.

20 DETAILED DESCRIPTION OF THE INVENTION

As mentioned above, patients treated with sulfonylureas gradually fail to respond to sulfonylurea treatment. It is generally accepted among those skilled in the art that this failure is due to exhaustion of β -cells which, accordingly,
25 are unable to excrete insulin in response to glucose stimulation. Also, it is generally accepted that the efficacy of sulfonylureas is limited by the capacity of β -cells to produce and excrete insulin. Accordingly, one would not expect any additional therapeutic advantage by treating NIDDM
30 patients with sulfonylureas and other agents stimulating β -cell function as well.

Our finding that NIDDM patients may advantageously be treated with GLP-1 related peptides in combination

with sulfonylureas or other oral hypoglycaemic agents is therefore, indeed, surprising. In fact, we have found that concomitant treatment with oral hypoglycaemic agents and GLP-1 related peptides results in a synergistic response by the NIDDM patients: treatment with oral hypoglycaemic agents and GLP-1 related peptides gives rise to a metabolic response greater than the sum of the responses of either agents when applied alone. Even in cases of sulfonylurea failures, the oral agents have been found to significantly enhance efficacy of GLP-1 related peptides.

Combined treatment with GLP-1 related peptides and oral hypoglycaemic agents is thus novel, therapeutically useful, and surprising. Unforeseen, therapeutic advantages can be gained by treating the NIDDM patients with both types of drugs.

Among the GLP-1 related peptides that can thus be used in the treatment of type 2 diabetes GLP-1(7-37) and GLP-1(7-36)amide are particularly advantageous, as they are identical to the naturally occurring hormones. Shorter peptides comprising part of the GLP-1(7-37) sequence or analogues of such shorter peptides or analogues of GLP-1(7-37) itself or functional derivatives of any of these can also be used to advantage, since pharmacodynamic and pharmacokinetic properties can be changed according to patients' demand by modifying the GLP-1 related fragment.

The GLP-1 related peptides can be administered by methods currently available according to the invention for administration of peptides. Nasal application is particularly advantageous from a patient compliance point of view. Details in this respect can be found in our copending Danish patent application No. DK 0364/92 relating to nasal administration of medicaments comprising GLP-1 related peptides which was filed simultaneously with the present application. The contents of said application is hereby incorporated in its entirety by reference. Administration by injection or infusion will be preferred in instances where a specific protracted plasma profile of the active peptide is

required, and oral administration is preferred in instances where extent and kinetics of absorption is not a critical issue.

The oral hypoglycaemic agent used according to the invention can be any oral agent exhibiting a glucose lowering effect. Among these agents, those acting on the ATP-dependent potassium channel of the β -cells are preferred such as glibenclamide, glipizide, gliclazide and AG-EE 623 ZW. The peptides according to the invention may also advantageously be applied in combination with other oral agents such as metformin and related compounds or glucosidase inhibitors as, for example, acarbose.

The features disclosed in the present description, examples and claims may, both separately and in any combination thereof, be material for realizing this invention in diverse forms thereof. The invention is further illustrated by the following examples which are not to be construed as limiting, but merely as an illustration of some preferred features of the invention.

20 **EXAMPLE 1**

Synergistic effect of GLP-1(7-36)amide and glibenclamide in NIDDM patients.

Assays

Blood samples were collected in plastic tubes containing EDTA (0.048 ml, 0.34 M) and Trasylol® (1000 IU Kallikrein inhibitor, obtained from Bayer, West Germany) and immediately placed on ice. The samples were centrifuged at 4°C and the plasma was stored at -20°C. Blood glucose was measured by a glucose oxidase method according to A.S. Hugget and D.A. Nixon, Lancet 2 (1957) 368-370. Plasma C-peptide concentrations were determined by radioimmunoassay (RIA) using a commercially available kit (Novo Research Institute, Denmark). Plasma glucagon concentrations were measured by RIA using antibody 30K as described by G.R. Faloon and R.H. Unger in B.M. Jaffe and Behrman, eds.

Methods of Hormone Radioimmunoassay, Academic Press, New York (1974) 317-330.

For further experimental details (e.g. on calculation of isoglycaemic meal-related insulin response, 5 IMIR), reference is made to M. Gutniak, C. Ørskov, J.J. Holst, B. Ahren and S. efendic, The New England Journal of Medicine 326 (29) (1992) 1316-1322, where a different experiment performed under similar conditions is described.

Methods

10 On four different days the effect of either injecting glibenclamide, 1 mg i.v., or infusing GLP-1(7-36)amide at a rate of 0.75 pmol per kilogram of body weight per minute or a combination thereof was studied in the same group of 6 insulin treated obese NIDDM patients (Body Mass 15 Index: 30.1 ± 2.4 kg/m²) and compared to administration of saline as control. Ordinary administration of insulin was stopped 24 hours before the administration of the test compounds or of the saline started and all subjects were fasted overnight. A Biostator (Miles, Diagnostic Division, 20 Elkhart, Ind.) was used for insulin administration in this period in order to normalize blood glucose levels before the administration of the test compounds was initiated and also to keep a normal postprandial blood glucose pattern 180 minutes following the ingestion of a standard test meal 25 comprising boiled potatoes, boiled beef, cooked carrots, a glass of milk containing 0.5% butterfat, and a slice of bread baked from a mixture of wheat and rye flours. In this meal, 28, 26, and 46% of the energy comes from protein, fat and carbohydrates, respectively. Administration of the test 30 compounds was performed (glibenclamide, saline) or initiated (GLP-1(7-36)amide, respectively, 30 minutes after normoglycaemia was achieved. The infusion of (GLP-1(7-36)amide was continued for 210 minutes. After 30 minutes (time zero), the subjects were given the test meal which was 35 consumed within 15 minutes. Blood samples were obtained at - 30, 0, 15, 30, 90, 120, 150 and 180 minutes.

Results

After the ingestion of the meal, meal-related C-peptide response, glucagon response and isoglycaemic meal-related insulin requirement (IMIR) was measured. The results are summarized in Table 1.

Table 1

	C-peptide response (pg/ml/210 min)	Glucagon response (pg/ml/210 min)	IMIR (U)
Control (saline)	7.4±3.6	269345±6299	17.4±2.8
GLP-1(7-36)amide	25±9.8	10451±5126	6.3±2.0
glibenclamide	105±53.9	*)	8.3±1.0
10 GLP-1(7-36)amide + glibenclamide	184±55.1	2526±4873	2.7±0.7

*) glibenclamide had no significant influence on glucagon release.

As indicated in the table, both GLP-1(7-36)amide and glibenclamide significantly increased meal-related C-peptide response ($p < 0.02$) and when administered in combination exerted a clear synergistic effect. GLP-1(7-36)amide suppressed the glucagon response ($p < 0.01$) while glibenclamide had no significant effect. However, in combination with GLP-1(7-36)amide the glucagon response was almost abolished. Finally, both glibenclamide and GLP-1(7-36)amide lowered IMIR and in combination IMIR was as low as 2.7±0.7.

In summary, this experiment demonstrates a strong synergistic effect of a combination of GLP-1(7-36)amide and glibenclamide.

EXAMPLE 2

Synergistic effect of GLP-1(7-36)amide and glibenclamide in NIDDM patients with secondary failure to sulfonylurea treatment.

5 Methods.

Eight patients with NIDDM and secondary failure to sulfonylurea treatment participated in the study (age 57.6 ± 2.7 years, body mass index 28.7 ± 1.5 kg/m², diabetes duration 7.6 ± 1.2 years, HbA_{1c} 5.8 ± 0.5). The diabetic
10 patients fulfilled the criteria for NIDDM and IDDM according to the USA National Diabetes Data Group. None of the patients had impaired renal function, automatic neuropathy, or proliferative retinopathy, and all had normal liver function. They were instructed to eat a standard diet for
15 diabetic patients at least 2 weeks before and during the study. The patients treated with sulfonylureas stopped their medication one week before the experiments. Those who were treated with insulin were instructed to stop the injections of NPH insulin 24 hours before the studies. Blood glucose
20 concentrations were controlled with subcutaneous injections of regular insulin.

All the subjects were studied after an overnight fast. At 07.30 h on the morning of each study, three cannulas were inserted. One cannula was placed in an
25 antecubital vein and was used to sample blood intermittently for hormone assays. It was flushed with saline after each sampling. A second cannula inserted retro-gradely in a dorsal hand vein was used for continuous monitoring of blood glucose concentrations. The venous blood was arterialized by
30 heating the forearm and hand in a thermoregulated sleeve (Kanthal Medical Heating AB, Stockholm,, Sweden) at 45°C. The third cannula was inserted in the contralateral antecubital vein and was used for all infusions. From approximately 08.00 hours, the patients were connected to a
35 Biostatator in order to normalize their blood glucose

concentrations. The algorithm of the Biostator was adjusted in order to normalize basal blood glucose levels. The target for blood glucose concentrations was 4-5 mmol/L. When the target was reached, the Biostator algorithm was changed to monitoring and the feedback insulin infusion was stopped. The experiments were started 30 minutes after normoglycemia was achieved, approximately 90 minutes after connection to the Biostator. An infusion of saline or 0.75 pmol/kg/min of GLP-1(7-36)amide (Peninsula Laboratories, St. Helens, Merseyside, England) then was started and continued for 210 minutes. In glibenclamide experiments an i.v. injection of 1 mg glibenclamide (Hoechst AG, Germany) was given at the same time point. These four studies were performed in a random order with 2-4 weeks elapsed between the experiments. At time 0 the subjects were given a standard lunch, as described in Example 1 which they ate within 15 minutes while sitting in bed. Blood samples were taken at FV, -60, -30, -15, 0, 15, 30, 90, 120, 150, and 180 minutes. Blood glucose was measured continuously.

Results.

In the basal state, the effect on blood glucose and C-peptide levels was monitored 45 minutes after administration of GLP-1(7-36)amide, glibenclamide or a combination thereof had started. The results are summarized in Table 2.

Table 2

	Blood glucose mmol/l	C-peptide pmol/l
Control (saline)	6.0 \pm 0.3	0.53 \pm 0.06
GLP-1(7-36)amide	5.1 \pm 0.4	0.63 \pm 0.1
glibenclamide	6.0 \pm 0.3	0.56 \pm 0.007
GLP-1(7-36)amide + glibenclamide	4.5 \pm 0.1	0.72 \pm 0.1

These results clearly demonstrates the synergistic effect of the two compounds as glibenclamide had no significant effect on its own while the effect of the combination of GLP-1(7-36)amide and glibenclamide, clearly, exceeded that of GLP-1(7-36)amide alone.

After the ingestion of the meal, the insulinogenic indices (integrated insulin/integrated glucose response) were calculated, again highlighting the synergistic effect of the two compounds, as shown in Table 3.

Table 3

	Insulinogenic index
Control (saline)	1.6 \pm 0.6
GLP-1(7-36)amide	21.0 \pm 7.2,
glibenclamide	10.6 \pm 2.8,
GLP-1(7-36)amide + glibenclamide	37.5 \pm 9

CLAIMS

1. Use of GLP-1(7-37), GLP-1(7-36)amide, or a pharmaceutically acceptable peptide containing a fragment of the GLP-1(7-37) sequence; or an analogue or a functional derivative of such a peptide for the preparation of a medicament for use in the treatment of diabetes in a regimen which additionally comprises treatment with an oral hypoglycaemic agent.
2. Use according to claim 1 of GLP-1(7-37) or GLP-1(7-36)amide.
3. Use according to claim 1 or 2 when the oral hypoglycaemic agent is a blocker of the ATP-dependent potassium channel on β -cells.
4. Use according to claim 1 or 2 when the oral hypoglycaemic agent is a sulfonylurea.
5. Use according to claim 1 or 2 when the oral hypoglycaemic agent is (S)-(+)-2-ethoxy-4-[2-[[3-methyl-1-[2-(1-piperidinyl)phenyl]butyl]amino]-2-oxoethyl]benzoic acid.
6. Use according to claim 1 or 2 when the oral hypoglycaemic agent is a biguanide.
7. Use according to claim 6 when the oral hypoglycaemic agent is metformin.
8. A method of treating diabetes which method comprises administering an effective amount of GLP-1(7-37), GLP-1(7-36)amide, or a pharmaceutically acceptable peptide containing a fragment of the GLP-1(7-37) sequence, or an analogue or a functional derivative of such a peptide to a patient in need of such a treatment in a regimen which additionally comprises treatment with an oral hypoglycaemic agent.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/DK 93/00099

A. CLASSIFICATION OF SUBJECT MATTER

IPC5: A61K 37/28

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC5: A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

BIOSIS, EMBASE, MEDLINE, WPI, CHEMICAL ABSTRACTS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Diabetes, vol 40(1991):Suppl. 1, page 237A, J.C. Parker et al: "Glucagon-like peptide 1(7-37) and glibenclamide stimulate insulin secretion by different glucose-dependent mechanisms". --	1-7
X	WO, A1, 8706941 (THE GENERAL HOSPITAL CORPORATION), 19 November 1987 (19.11.87) --	1-7
X	WO, A1, 9011296 (THE GENERAL HOSPITAL CORPORATION), 4 October 1990 (04.10.90) --	1-7

☒ Further documents are listed in the continuation of Box C.☒ See patent family annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

† later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

23 June 1993

Date of mailing of the international search report

28 -06- 1993

Name and mailing address of the ISA/
Swedish Patent Office
Box 5055, S-102 42 STOCKHOLM
Facsimile No. +46 8 666 02 86

Authorized officer

Elisabeth Carlborg
Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT

International application No.

PCT/DK 93/00099

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO, A1, 9111457 (BUCKLEY, DOUGLAS I. ET AL), 8 August 1991 (08.08.91) ----- --	1-7

INTERNATIONAL SEARCH REPORT

International application No.

PCT/DK 93/00099

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 8
because they relate to subject matter not required to be searched by this Authority, namely:
See PCT Rule 39.1(iv): Methods for treatment of the human or animal body by surgery or therapy, as well as diagnostic methods.
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT
Information on patent family members

28/05/93

International application No.

PCT/DK 93/00099

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A1- 8706941	19/11/87	EP-A- 0305387	08/03/89
WO-A1- 11296	04/10/90	NONE	
WO-A1- 9111457	08/08/91	EP-A- 0512042	11/11/92